arXiv:gr-qc/0407090v4 11 Jan 2006

Black Holes Conserve Information in Curved-Space Quantum Field Theory

Christoph Adami[1,2,3] and Greg L. Ver Steeg[2]
[1] *Jet Propulsion Laboratory 126-347,*
California Institute of Technology, Pasadena, CA 91109
[2] *California Institute of Technology 103-33,*
Pasadena, CA 91125
[3] *Keck Graduate Institute, Claremont, CA 91711*

(Dated: February 7, 2008)

The fate of classical information incident on a quantum black hole has been the subject of an ongoing controversy in theoretical physics, because a calculation within the framework of semi-classical curved-space quantum field theory appears to show that the incident information is irretrievably lost, in contradiction to time-honored principles such as time-reversibility and unitarity. Here, we show within the same formalism embedded in quantum communication theory that signaling from past to future infinity in the presence of a Schwarzschild black hole can occur with arbitrary accuracy, and thus that information is not lost in black hole dynamics. The calculation relies on a treatment that is manifestly unitary from the outset, where probability conservation is guaranteed because black holes stimulate the emission of radiation in response to infalling matter. This radiation is *non-thermal*, and contains all of the information about the infalling matter, while Hawking radiation contains none of it.

PACS numbers: 04.70Dy,03.67-a,03.65.Ud

Black hole evaporation poses a serious challenge to theoretical physics because it is a problem at the intersection of general relativity and quantum mechanics, precisely where our understanding is the weakest. From the moment Hawking discovered the eponymous radiation effect [1], many researchers have concluded that black holes introduce an intrinsic *irreversibility* into the laws of physics, together with an intrinsic *unpredictability* [2]. The reasons for such a radical departure from conventional physics at first seem incontrovertible. Because the only identifiers (quantum numbers) of a black hole that can be measured are its mass, charge, and angular momentum, information not encoded in these quantum numbers would be lost after the particle disappeared behind the horizon, making the dynamics strictly irreversible. This implies that, from the point of view of communication theory, signaling between a sender that encodes information into a beam of particles aimed at the horizon, for example, and a receiver at rest outside the horizon at future infinity, would be fundamentally impossible.

If information is trapped within a black hole, we are faced with another paradox, known as the *coherence problem* [3]. Hawking radiation, while reducing the mass of the black hole, results in a highly entangled quantum state whose support is the joint Hilbert space spanned by the horizon and future infinity. While this state can be pure, the quantum state at future infinity (averaged over the horizon) is naturally mixed. If black hole evaporation ultimately results in the disappearance of the horizon, what happens to the quantum state at this point? Does it remain mixed, or does it magically return to a pure state on future infinity? Standard arguments suggest that the latter cannot happen if information remains trapped behind the horizon, because black holes just before complete evaporation would be too small to encode all of it. And even if they could, this much entropy could not be radiated away in a finite amount of time. Technically speaking, the coherence problem is different from the information paradox because it addresses the quantum state *after* black hole evaporation, while the information paradox concerns the possibility of signaling in the presence of a horizon. However, the two problems are intimately related, because if information does not “accumulate” behind the horizon but instead remains outside (as we suggest), then the coherence problem appears much less severe.

Here, we address the information paradox, or more precisely, the impossibility of signaling in the presence of a black hole event horizon, by showing that the information transmission capacity of the black hole channel is non-zero, and that therefore classical signal states can be decoded with arbitrary accuracy by an observer suspended at future infinity outside the black hole horizon. We derive the black hole channel capacity within standard curved-space quantum field theory, that is in the semiclassical approximation where the back-reaction of the quantum dynamics on the metric field is ignored. Signaling is made possible by the *stimulated emission* of radiation that must accompany absorption. The well known *spontaneous emission* (Hawking radiation) turns out to provide the noise of the black hole channel, noise from which no information can be extracted. In the following, we first calculate the joint quantum state of black hole and radiation when late-time (signaling) particles accrete, then use standard quantum channel theory (see, e.g., [4]) to calculate the capacity of the channel.

In units where $\hbar = c = G = k = 1$, a non-charged, non-rotating black hole has an entropy given by $S_{\rm BH} = 4\pi M^2$, where M is the mass of the black hole, and a temperature $T = (8\pi M)^{-1}$, the Hawking temperature. Black holes are formed in stellar collapse of stars of sufficient mass, and can subsequently accrete particles (see Fig. 1a). Through the process of virtual pair formation near the event horizon, with one of the pair's particles disappearing behind the event horizon and the other going off to future infinity (the time-slice $\mathcal{I}^+$ in Fig. 1), a black hole loses mass by providing it to the virtual pair, which, in turn, goes on mass shell. The problem of loss of predictability is easily seen in Fig. 1a, where different trajectories of particles accreting onto the black hole are outlined, and arbitrarily labelled. If the identity of these labels is lost behind the horizon, we are faced with a many-to-one mapping and concurrent loss of predictability (even if this information is released at a later time) because the future state of the black hole cannot be ascertained given the labels of the accreting particles. If the labels are destroyed in the singularity, then we would be forced to assume that, to make matters worse, the laws of physics allow non-unitary, and thus by extension probability non-conserving, dynamics.



FIG. 1: (a) Penrose diagram of the spacetime of a black hole, with accretion of arbitrarily labelled particles from past infinity ($\mathcal{I}^-$). Once the particles cross the event horizon into region II, they are indistinguishable (leading to a loss of information) unless they leave a signature outside (in region I) via stimulated emission (labelled α', β', γ'). (b) Modes a, b, c and A are concentrated in a region of null infinity indicated by the letter (note that a and b actually overlap on $\mathcal{I}^-$).

To introduce our notation, we begin by recapitulating the standard quantum theory of complex scalar fields in a gravitational background due to Hawking [1]. The presence of a black hole's event horizon introduces a complication to quantum field theory not encountered in flat-space: different vacuum states can be defined that are inequivalent in terms of their particle content. The so-called *Unruh* vacuum is non-trivial due to the separation of Schwarzschild space-time into two regions, generally depicted as the "outside" and "inside" region, or region I and region II, respectively (Fig. 1). In the standard treatment, the operators that annihilate the curved-space vacuum, A_k and B_k (A_k annihilates the "outside" vacuum while B_k annihilates "inside"), are related to the flat space operators a_k and b_k by a Bogoliubov transformation [5], so for example

$$A_k \;=\; e^{-iH}a_k e^{iH} = \alpha a_k - \beta b^\dagger_{-k}\ . \tag{1}$$

where $a^\dagger_k$ creates a particle of mode k while a_{-k} annihilates an antiparticle of the same mode. The transformation provides a mapping from the trivial *Boulware* vacuum $|0\rangle_g = |0\rangle_a|0\rangle_b$ (the Boulware vacuum equals the standard Minkowski vacuum far from the black hole) to the Unruh vacuum $|0\rangle_U$, using a unitary operator $U = e^{-iH}$ [5]

$$|0\rangle_U = e^{-iH}|0\rangle_g\ . \tag{2}$$

The Hermitian operator

$$H = ig(a^\dagger_k b^\dagger_{-k} - a_k b_{-k} + k \to -k) \tag{3}$$

describes the entanglement of particles and antiparticles outside and inside the horizon (for mode k). In terms of the interaction strength g, it follows that

$$\alpha^2 \;=\; \cosh^2 g = \frac{1}{1-e^{-\omega/T}} \tag{4}$$

$$\beta^2 \;=\; \sinh^2 g = \frac{1}{e^{\omega/T}-1}\ , \tag{5}$$

where $\omega = |k|$ is the frequency associated with mode k and T is the Hawking temperature. The relationship between the coefficients α and β and the Hawking temperature T is enforced by an analyticity condition on the solution to the free-field equations, as usual [1].

According to Eq. (2), U acts as an S-matrix mapping the in-states on past infinity ($\mathcal{I}^-$ in Fig. 1) into the out-states on future infinity ($\mathcal{I}^+$ and the horizon, in Fig. 1). As is well known, the Bogoliubov transformation (1) leads to a thermal density matrix for both particles and antiparticles when tracing $\rho = |0\rangle_U\langle 0|$ over the complementary sector, e.g.,

$$\rho_k = (1-e^{-\omega/T})\sum_{n_k=0}^{\infty} e^{-n_k\omega/T}|n_k\rangle_g\langle n_k|\ . \tag{6}$$

The mean number of particles emitted in region I (mode a) is also easily calculated from (1) and (2) as

$$\langle m\rangle = {}_U\langle 0|a^\dagger_k a_k|0\rangle_U = {}_g\langle 0|A^\dagger_k A_k|0\rangle_g = \beta^2\ , \tag{7}$$

which is the famous Planck distribution of Hawking radiation. From (6) we can calculate the von Neumann entropy of frequency mode k of the inside or outside region as

$$S(\rho_k) = \frac{\omega/T}{e^{\omega/T}-1} + \log(1-e^{-\omega/T}) \ . \tag{8}$$

This entropy is completely thermal, which implies that it does not depend on the distribution of matter that formed the black hole, or on the identity of particles absorbed by the black hole at late times (our signaling states). No information can be extracted from it.

However, as has been pointed out several times in the past [6, 7, 8, 9, 10, 11, 12], the standard treatment that we have repeated above actually ignores an important physical process that leads to a crucial modification of the dynamics, and in hindsight ensures the possibility of signaling. This process is *stimulated emission*, which must be present when absorption and spontaneous emission can occur. In fact, as is well known since Einstein's seminal work [13], the probability for stimulated emission is directly proportional to that of spontaneous emission. Just as the vacuum emits pairs of particles of all types, momenta, and particle numbers spontaneously (here, with probability β^2), it also reacts to *infalling*, that is, absorbed, radiation or particles, by emitting instead an exact *copy* of the absorbed radiation or particle. Note that the concomitantly emitted Hawking radiation prevents violation of the no-cloning theorem (see [14]).

Stimulated emission due to particles present just before formation of the black hole (early-time modes) can be described within the formalism just introduced [6, 8, 10], but does not shed light on the fate of information incident on the black hole after its formation, by *late-time* modes. Such modes (which we denote by c) are strongly blueshifted with respect to the early-time modes a and b that are the positive and negative-frequency parts of matter and radiation that propagate just outside and just inside the horizon introduced earlier, and therefore commute with them [9]. These modes are indicated schematically in Fig. 1b.

The Bogoliubov transformation connecting the outgoing mode A_k to a_k and the late-time mode c_k can be taken as:

$$A_k \ = \ e^{-iH}a_k e^{iH} = \alpha a_k - \beta b^\dagger_{-k} - \gamma c_k \ , \tag{9}$$

with $\alpha^2 - \beta^2 + \gamma^2 = 1$ due to unitarity.

The additional term γc_k turns out to be necessary for a consistent treatment of black hole thermodynamics [9], and can be accounted for by using an effective Hamiltonian with an extra term (compared to Eq. 3) that describes scattering of late-time modes c_k by the static black hole horizon, with an interaction strength g':

$$H = ig(a^\dagger_k b^\dagger_{-k} - a_k b_{-k}) + ig'(a^\dagger_k c_k - a_k c^\dagger_k) + k \to -k \ . \tag{10}$$

Such an interaction is entirely natural within quantum field theory and is completely analogous to Hamiltonians used in quantum optics to describe transformations in linear optics [15]. In fact, the first term (the standard Hawking term) corresponds to the Hamiltonian of an *active* optical element leading to state squeezing with strength g, while the second one describes a *passive* element: a simple beam splitter with interaction strength g'. We will discuss the magnitude of this coupling constant in relation to the early-time interaction strength g below, but can assume that it will not be very different as they are intrinsically the same type of particles.

Using (10) in (9), the Bogoliubov coefficients α, β, and γ can be expressed in terms of g and g' as

$$\alpha^2 \ = \ \cos^2(g'w) \ , \tag{11}$$

$$\beta^2 \ = \ (\frac{g}{g'})^2 \frac{\sin^2(g'w)}{w^2} \ , \tag{12}$$

$$\gamma^2 \ = \ \frac{\sin^2(g'w)}{w^2} \ , \tag{13}$$

where $w^2 = 1 - (g/g')^2$.

We can now calculate the reduced density matrix of outgoing radiation in region I when no particles are incident (notation $k|0$) just as before, by tracing the full density matrix $|0\rangle_U\langle 0|$ over region II (containing modes b and c), and find

$$\rho_{k|0} = \frac{1}{1+\beta^2}\sum_{n_k=0}^{\infty}\left(\frac{\beta^2}{1+\beta^2}\right)^{n_k} |n_k\rangle\langle n_k| \ . \tag{14}$$

If we identify β^2 with the expression $\frac{\Gamma}{e^{\omega/T}-1}$, (14) is precisely the density matrix of spontaneous emission from a black hole with absorptivity Γ, previously obtained using maximum entropy methods by Bekenstein [16]. It reduces to (6) in the limit $\Gamma \to 1$. Thus, the scattering term in (10) simply provides a microscopic model for late-time particle scattering. Using the aforementioned expression for β^2 along with the relation $\beta^2 = e^{-\omega/T}\alpha^2$ (which is due to the standard analyticity argument for the solution to the field equations that carries through as before) we also identify

$$\alpha^2 = \frac{\Gamma}{1-e^{-\omega/T}} \ , \tag{15}$$

and $\gamma^2 = 1 - \Gamma$, owing to $\alpha^2 - \beta^2 + \gamma^2 = 1$.

Consider now the outgoing state $|n_k\rangle_U$ when n_k late-time particles are incoming, constructed as before via

$$|n_k\rangle_U = e^{-iH}|n_k\rangle_g \ , \tag{16}$$

where $|n_k\rangle_g = |0\rangle_a|0\rangle_b|n_k\rangle_c$, i.e., the state with n_k incoming particles in mode c on $\mathcal{I}^-$. From Eq. (9), we immediately obtain the number of particles emitted into mode a if n_k were incident in mode c, since

$${}_U\langle n_k|a^\dagger_k a_k|n_k\rangle_U = {}_g\langle n_k|A^\dagger_k A_k|n_k\rangle_g = \gamma^2 n_k + \beta^2 \ . \tag{17}$$

This shows that, in addition to the standard spontaneous emission term β^2, detectors at infinity record $\gamma^2 n_k$ particles at infinity. The latter particles comprise two contributions: $(1-\alpha^2)n_k$ due to elastic scattering with a single quantum absorption probability α^2, and $\beta^2 n_k$ particles due to stimulated emission.

The density matrix of outgoing radiation in region I when n_k particles are incident (notation $k|n$) can be calculated from $|n_k\rangle_U\langle n_k|$ by repeated application of the disentangling theorems for SU(2) and SU(1,1), and tracing over the degrees of freedom of region II (modes b and c). Because no antiparticles are sent in, the antiparticle part of the density matrix factorizes and we can write $\rho_a^{(n)} = \rho_{k|n} \otimes \rho_{-k|0}$ with

$$\rho_{k|n} = \sum_{m_k=0}^{\infty} p(m_k|n_k)|m_k\rangle\langle m_k| \ . \tag{18}$$

The expression for $p(m|n)$, the probability to detect m outgoing particles at $\mathcal{I}^+$ if n were incident on the black hole is complicated, but agrees exactly with that obtained by Bekenstein and Meisels [7] using maximum entropy methods, and by Panangaden and Wald [8] in quantum field theory. Note that because incoming late-time particles stimulate the emission of *anti*-particles behind the horizon, the stimulated emission process conserves all quantum numbers.

We can use our expression for $p(m|n)$ to put limits on the scattering strength g'. For example, the probability that no particle is found outside the horizon while a single quantum is incident is given by

$$p(0|1) = \frac{\alpha^2}{(1+\beta^2)^2} = \frac{\alpha^2}{(1+\alpha^2 e^{-\omega/T})^2} \ , \tag{19}$$

which tends to the single-quantum absorption probability α^2 in the zero-temperature limit. Thus, while g sets the temperature of the black hole as in the standard formalism, g' in turn sets the reflectivity of the grey body. Because $0 \leq p(0|1) \leq 1$, we can deduce that $0 \leq \alpha^2 \leq 1$, which in turn implies $g' \geq g$ or else $g' = 0$ (see Eq. 11). The latter solution corresponds to the familiar Hawking Hamiltonian (3) where $\Gamma = \alpha^2 = 1$, but cannot account for a thermodynamically consistent picture that includes scattering from the potential barrier surrounding the black hole ($\Gamma < 1$), as pointed out by Bekenstein and Meisels [7].

A preferred choice is $g' = g$, as this limit describes perfect absorption ($\alpha^2 = 1$) where we find

$$p(m|n) = \frac{e^{-m\omega/T}}{(1+e^{-\omega/T})^{n+m+1}} \ . \tag{20}$$

But even in this extreme case, the classical (effective) absorption probability $\Gamma = \alpha^2(1-e^{-\omega/T})$ is strictly smaller than unity due to the stimulated emission effect, thus allowing for the preservation of information.

To show this, let us cast black-hole dynamics into the language of quantum information theory. We can follow the fate of information interacting with a black hole by using a *preparer* to encode information into late-time quantum states that are then sent into the event horizon. For example, we can imagine a preparer X who sends packets of n particles with probability $p(n)$. The internal state of the preparer can be described by the density matrix $\rho_X = \sum_n p(n)|n\rangle\langle n|$, with entropy $S(\rho_X) = H[p] = -\sum_n p(n)\log p(n)$. After the particles interact with the black hole, our preparer is now correlated with it because the final state is now the density matrix

$$\rho_{\mathrm{I,II},X} = \sum_n p(n)|n\rangle_U\langle n| \otimes |n\rangle_X\langle n| \ , \tag{21}$$

Tracing over the black-hole interior (region II), we obtain the joint density matrix of the radiation field in region I and the preparer X:

$$\rho_{\mathrm{I},X} \ = \ \sum_n p(n)\rho_{k|n} \otimes |n\rangle_X\langle n| \tag{22}$$

with entropy

$$S(\rho_{\mathrm{I},X}) = H[p] + \sum_n p(n)S(\rho_{k|n}) \tag{23}$$

owing to the block-diagonal form of Eq. (22). The mutual entropy between the radiation field at future infinity and the preparer is then simply given by

$$\begin{aligned} H(X:\mathrm{I}) \ &= \ S(\rho_{\mathrm{I}}) + S(\rho_X) - S(\rho_{\mathrm{I},X}) \\ &= \ S\left(\sum_n p(n)\rho_{k|n}\right) - \sum_n p(n)S(\rho_{k|n}) \ , \end{aligned} \tag{24}$$

which is known as the Holevo bound [17]. The latter constitutes the maximum amount of classical information that can be extracted from a quantum measurement, and its maximum (over the probability distribution of signal states) turns out to be the capacity of a quantum channel to transmit classical information [18].

As a simple example, consider a binary channel where the preparer either sends no particle (with probability $1-p$) or a packet of n particles (with probability p) into the black hole. The resulting density matrix $\rho_{k|n}$ of outgoing particles in region I is diagonal in the number basis. For example, for $n = 1$

$$\rho_{k|1} \ = \ \frac{\alpha^2}{(1+\beta^2)^2} \sum_{m_k=0}^{\infty} \left(\frac{\beta^2}{1+\beta^2}\right)^{m_k} (1+m_k\frac{\gamma^2}{\alpha^2\beta^2})|m_k\rangle\langle m_k| \tag{25}$$

is clearly non-thermal, except in the unphysical Hawking limit $g' = 0$. This implies that the channel's capacity



FIG. 2: (a) Capacity of the binary black hole channel ($n = 1$) for different absorption probabilities α^2, as a function of ω/T. The capacity of the noiseless channel is $\chi = 1$. (b) Capacity for anti-particle/particle encoding (solid line) and particle/vacuum encoding, as a function of the packet size n.

$\chi = \max_p H(X : \text{I})$ never vanishes (as $g' \geq g$), and information can be recovered with perfect accuracy (see Fig. 2).

Of course, absorption of particles by the black hole dampens the signal, so that the capacity declines with α^2. However, even in the worst case (full absorption, $\alpha^2 = 1$) the capacity remains positive: $\chi \sim e^{-\omega/T}$, and information can still be recovered with arbitrary accuracy, using standard methods of error correction. Note furthermore that the information loss in the absence of coding occurs *outside* of the event horizon (via the mixing of stimulated signal modes with Hawking noise). We can see this by encoding our logical bit into particle states that are more easily distinguished from Hawking radiation. For example, sending bunches of n particles with $n > 1$ to signal a "1" while using the vacuum as the "0-state" yields a much larger capacity (dotted line in Fig. 2b). Using anti-particle bunches of size n to signal "0" increases this capacity again (solid line), as these are much easier to detect than the absence of particles.

In conclusion, we found that a consistent treatment of black hole dynamics requires the presence of stimulated radiation outside the event horizon beyond the usual Hawking radiation. The radiation field in this region is non-thermal, while accretion and evaporation described by the Hamiltonian (10) is unitary. This guarantees that any change in the entropy of the inside region is exactly balanced by a commensurate change in the outside. The information dynamics of black holes turn out to be those of a standard noisy quantum channel that is used to transmit classical information, where Hawking radiation provides the noise source. This calculation does not address the fate of the black hole *after* complete evaporation directly, but strongly suggests that nothing surprising needs to happen there, because the information about the initial state that formed the black hole, as well as about any late-time particles accreting on it, is available *outside* the event horizon in the form of stimulated emission.

Acknowledgements We are grateful to N.J. Cerf and C.O. Wilke for crucial discussions and comments on the manuscript. We also thank J.P. Dowling, D. Gottesman, G.M. Hockney, P. Kok, H. Lee, and U. Yurtsever for discussions. This work was carried out in part at the Jet Propulsion Laboratory (California Institute of Technology) under a contract with NASA, with support from the Army Research Office's grant # DAAD19-03-1-0207.

[1] Hawking, S. W. Particle creation by black holes. *Commun. Math. Phys.* **43**, 199–220 (1975).

[2] Hawking, S. W. Breakdown of predictability in gravitational collapse. *Phys. Rev. D* **14**, 2460 (1976).

[3] Wald, R. M. *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics* (University of Chicago Press, Chicago, IL, 1994).

[4] Preskill, J. Quantum Computation and Information. Phy 219 Lecture Notes (Caltech) (1998).

[5] Birrell, N. D. & Davies, P. C. W. *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).

[6] Wald, R. M. Stimulated-emission effects in particle creation near black holes. *Phys. Rev. D* **13**, 3176–3182 (1976).

[7] Bekenstein, J. D. & Meisels, A. Einstein a and b coefficients for a black hole. *Phys. Rev. D* **15**, 2775 (1977).

[8] Panangaden, P. & Wald, R. M. Probability distribution for radiation from a black hole in the presence of incoming radiation. *Phys. Rev. D* **16**, 929 (1977).

[9] Sorkin, R. A simplified derivation of stimulated emission by black holes. *Class. Quant. Grav.* **5**, L149–L155 (1987).

[10] Audretsch, J. & Müller, R. Amplification of black-hole hawking radiation by stimulated emission. *Phys. Rev. D* **45**, 513 (1992).

[11] Schiffer, M. Is it possible to recover information from the black-hole radiation? *Phys. Rev. D* **48**, 1652 (1993).

[12] Müller, R. & Lousto, C. Recovery of information from black hole radiation by considering stimulated emission. *Phys. Rev. D* **49**, 1922–1928 (1994).

[13] Einstein, A. Zur quantentheorie der strahlung. *Physik. Zeitschr.* **18**, 121–128 (1917).

[14] Simon, C., Weihs, G. & Zeilinger, A. Optimal quantum cloning via stimulated emission. *Phys. Rev. Lett.* **84**, 2993–2996 (2000).

[15] Leonhardt, U. Quantum physics od simple optical instruments. *Rep. Progr. Phys.* **66**, 1207 (2003).

[16] Bekenstein, J. Statistical black-hole thermodynamics. *Phys. Rev. D* **12**, 3077–3085 (1975).

[17] Holevo, A. Some estimates of the information transmitted by quantum communication channel. *Probl. Inf. Transm. (USSR)* **9**, 177 (1973).

[18] Holevo, A. The capacity of quantum channel with general signal states. *IEEE Trans. Info. Theor.* **44**, 269–273

(1998).